arXiv:1301.1895v1 [gr-qc] 9 Jan 2013

Effect of eccentricity on binary neutron star searches in Advanced LIGO

E. A. Huerta and Duncan A. Brown
Department of Physics, Syracuse University, Syracuse, NY 13244, USA and
Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106.

(Dated: January 10, 2013)

Binary neutron stars (BNSs) are the primary source of gravitational waves for the Laser Interferometer Gravitational-wave Observatory (LIGO) and its international partners Virgo and KAGRA. Current BNS searches target field binaries whose orbits will have circularized by radiation reaction before their gravitational waves enter the Advanced LIGO sensitive band at 15 Hz. It has been suggested that a population of BNSs may form by n-body interactions near supermassive black holes or in globular clusters and that these systems may have non-negligible eccentricity in the Advanced LIGO band. We show that for BNS systems with total mass of $2.4\,M_\odot$ ($6\,M_\odot$), the effect of eccentricity $e \lesssim 0.02$ (0.05) is negligible and a circular search is effectual for these binaries. For eccentricities up to $e = 0.4$, we investigate the selection bias caused by neglecting eccentricity in BNS searches. If such high eccentricity systems exist, searches that specifically target eccentric binaries will be needed in Advanced LIGO and Virgo.

I. INTRODUCTION

Binary neutron star (BNS) mergers are expected to be rich sources of gravitational waves (GWs) [1] and the progenitors of short gamma-ray bursts (SGRBs) [2], among other electromagnetic counterparts [3]. These sources may be detected by advanced ground-based GW observatories [4–6] within a horizon distance of 445 Mpc [7]. Observations of galactic double NS systems containing pulsars, which formed through binary evolution in the disk of the galaxy, suggest that five of these systems are expected to merge through GW radiation within a Hubble time [8, 9]. Observed field BNSs have orbital periods of a few hours, merger times $\sim 10^{2-3}$ Myr, low-kick velocities $\lesssim 50\,\mathrm{km\,s^{-1}}$, and are expected to enter the frequency band of ground-based GW detectors with negligible residual eccentricities [8]. For example the binary pulsar PSR 1913+16 is expected to have an eccentricity $e \sim 10^{-6}$ when its gravitational-wave frequency is 15Hz [8, 10, 11]. Population synthesis models constrained by the double NS systems in the field predict a binary coalescence rate in Advanced LIGO (aLIGO) of 0.4–400 per year, with a likely value of 40 per year [7].

In addition to the observed field binaries, observations by Swift have detected a population of SGRBs in or near elliptical galaxies [12]. Some of these SGRBs have spatial offsets that cannot be attributed to the ejection of the progenitor compact binary from the host galaxy by the kick imparted to the system through the supernova explosions that create its components [13]. The necessary kick velocities for the progenitor compact binary to remain bound to the elliptical host galaxy and at the same time have its apogalacticon in the halo are very different to those observed in field BNSs [14]. In [13], Grindlay et al. propose that SGRBs observed in or near ellipticals may be produced by the mergers of BNSs formed by dynamical capture in core-collapsed globular clusters. In these dense stellar environments, n-body interactions may lead to the formation of binaries that have sizable eccentricities at merger [15]. Based on the globular-cluster population of elliptical galaxies, and scaling from the BNS observed in the globular cluster M15 [16], Grindlay et al. showed that around $10-30\%$ of SGRBs may be generated from BNSs in globular clusters, although the expected detection event rate of BNSs in globulars for ground-based antennas is a factor ~ 200 smaller than the expected rate for field BNSs [7, 17].

Dynamical interactions in galactic nuclei may also lead to the formation of eccentric BNSs [18, 19]. BNSs formed near galactic nuclei may enter the LIGO band before the maximum eccentricity expected from Kozai oscillations is reached. After multiple periodic oscillations in the orbital elements, and before gravitational radiation becomes important, these BNSs may enter the LIGO band with high eccentricities [19]. Nonetheless, the contribution of these type of sources to the BNS coalesce rates is likely to be small, namely: 10, 1 and 0.1% of the low, realistic and high estimates in [7]. In addition to BNS mergers, stellar-mass black holes (BHs) are expected to segregate and form steep cusps around supermassive BHs in galactic nuclei. Emission of gravitational radiation during encounters among these self-segregated BHs will drive the formation and coalescence of binary BHs that aLIGO may detect with a rate $\sim 1 - 10^2\mathrm{yr}^{-1}$. This optimistic estimate depends on the initial mass function of stars in galactic nuclei and the mass of the heavier BHs. These systems will be distinguishable from other binaries because more than 90% of them are expected to have significant eccentricity ($e > 0.9$) when they enter LIGO band [18]. The studies quoted previously do not take into account natal kicks. In [20], it is shown that if a BNS system receives a supernova kick which is similar in magnitude to its circular speed, then the likelihood of undergoing an extremely fast coalescence is very high. Indeed, for BNS systems that experience planar kicks, about 24% of them coalesce ten thousand times faster than a BNS system that stays in a circular orbit immediately after the supernovae. Pulsar surveys may not detect these events because current search algorithms are not equipped to cope the strong Doppler smearing effects present in these systems [20]. A pulsar can be detected if its lifespan is of the order of 10 Myrs and so highly eccentric systems which are driven to coalescence within 1Myr or less constitute a currently ‘missing’ BNS population that aLIGO may detect.

Motivated by the possibility of an additional population of eccentric binaries, we revise the study carried out in [21] to asses the importance of including eccentricity in BNS searches in the advanced detector era. Eccentric BNSs may not contribute significantly to the total number of BNS events that may be detected, but this study quantifies the selection bias introduced by ignoring eccentricity in aLIGO searches. Based on results of radio surveys that have confirmed the existence of pulsars with masses heavier than the canonical value of $1.35M_\odot$ [22], we carry out a large Monte Carlo simulation to address the efficacy of current search pipelines to detect eccentric BNS events in the mass range $1M_\odot - 3M_\odot$, making use of the Zero Detuned High Power (ZDHP) noise curve for aLIGO [23]. To simulate the eccentric BNS signals, we use the model introduced in [24], which consists of a post-Newtonian (PN) model that was calibrated by comparison to a numerical relativity simulation of an eccentric, equal-mass binary black hole. This eccentric PN model combines the 3PN conservative quasi-Keplearian orbit equations with the 2PN evolution of the orbital elements to construct an adiabatic waveform that best matches the numerical relativity simulation. We use the implementation of this model, described in [21], with the addition of an adaptive mesh-refinement method and an accelerated integrator. These improvements to the model were necessary to carry out a large scale Monte Carlo simulation to study the efficacy with which GW searches for circular binaries may be able to detect BNS signals with non-negligible eccentricity in the observational band of advanced detectors. In Section II we describe the method used to construct a template bank of non-eccentric BNS waveforms. Section III presents the results of our investigations. Finally, Section IV presents a summary of our findings and future directions of work.

II. BANK SIMULATION STUDIES

In this Section, we discuss the efficacy with which a template bank [25, 26] of non-eccentric waveforms can recover GW signals of eccentric BNS systems. The Fourier transform of a gravitational-wave signal $h(t)$ is given by

$$\tilde{h}(f) = \int_{-\infty}^{\infty} e^{-2i\pi f t} h(t) \mathrm{d}t \,. \tag{1}$$

The noise-weighted inner product between two signals h_1 and h_2 is given by

$$(h_1|h_2) = 2\int_{f_{\min}}^{\infty} \frac{\tilde{h}_1^*(f)\tilde{h}_2(f) + \tilde{h}_1(f)\tilde{h}_2^*(f)}{S_n(f)} \mathrm{d}f \,, \tag{2}$$

where $S_n(f)$ is the power spectral density of the detector noise. We use the form of the aLIGO ZDHP noise spectrum and set the lower limit of the integral in Eq. (2) to $f_{\min} = 15$Hz. From this we can construct the matched-filter signal-to-noise ratio [27]

$$\rho = \frac{(s|h)}{\sqrt{(h|h)}} \,, \tag{3}$$

where $s = n(+h)$ is the output of the detector, and n is the detector noise. The normalized overlap between any two waveforms h_1, h_2 is

$$\left(\hat{h}_1, \hat{h}_2\right) = \frac{(h_1|h_2)}{\sqrt{(h_1|h_2)\,(h_1|h_2)}} \,. \tag{4}$$

The time of coalesce t_c and phase of coalescence ϕ_c are maximized over in both the SNR and the overlap [28]. The maximized overlap

$$\mathcal{O}\left(h_1, h_2\right) = \max_{t_c,\,\phi_c} \left(\hat{h}_1, \hat{h}_2\right) \,, \tag{5}$$

gives the loss in signal-to-noise ratio incurred due to mismatches in the waveforms [29]. In this paper we compute the effectualness of non-eccentric templates to recover eccentric systems. To do this, we construct a ‘bank’ of templates which cover the space of circular BNS signals. The templates are place on a hexagonal lattice using the method introduced in [30, 31], so that the loss in signal-to-noise ratio between a signal and the nearest template is no more than 3%. The metric used to place the template grid is constructed using TaylorF2 inspiral templates that include corrections to the orbital phase evolution at 2PN order. The waveform family we will use to create the template bank waveforms is Taylor T4 [32, 33], which includes corrections to the phase of the waveform to 3.5PN order [34–36]. We have confirmed that this bank is effectual for TaylorT4 waveforms before considering the effect of eccentricity. We then compute the correlation or overlap between a simulated eccentric GW signal (h^e) and the best fitting non-eccentric template (h_b^T) by maximizing the overlap over the templates in the bank. This is known as the Fitting Factor ($\mathcal{FF}$) [29], and is defined as

$$\mathcal{FF} = \max_{b \in \mathrm{bank}} \mathcal{O}(h^e, h_b^T) \,. \tag{6}$$

In the next section we present the results of this fitting factor investigation using the eccentric x-model waveforms as signals.

III. RESULTS

The Monte Carlo simulation includes 40,000 points uniformly distributed in individual component mass $1M_\odot < m_{1,\,2} < 3M_\odot$, and with a uniform distribution of eccentricity in the range $e \in [0, 0.4]$. The template waveforms are generated from an initial gravitational wave frequency of 15Hz. In order to avoid edge effects, we generate the eccentric waveforms using an initial gravitational wave frequency of 14Hz.

The left panel of Figure 1 presents a cumulative histogram of the injected eccentric signals with $\mathcal{FF}$ less than the value associated on the x-axis. There are two intrinsic parameters that determine the effectualness of the circular bank to recover eccentric signals. The right panel of Figure 1 shows that the $\mathcal{FF}$ decreases for increasing values of eccentricity, as expected. The right panel of Figure 1 also shows the correlation



FIG. 1. The left panel is a cumulative histogram which indicates the fraction of points in the mass-pace region where the bank of non-eccentric waveforms has a value of Fitting Factor ($\mathcal{FF}$) less than the associated value on the x-axis. The right panel shows the $\mathcal{FF}$ as a function of the initial value of the eccentricity of the binary system.

between the $\mathcal{FF}$ and eccentricity for systems with component masses $m_{1,2} \in [1M_\odot, 3M_\odot]$, and suggests that some systems can be recovered with $\mathcal{FF} \gtrsim 0.95$ in an eccentricity range $e \in [0.02, 0.05]$.

Figure 2 shows that for BNSs the $\mathcal{FF}$ depends primarily on the initial value of the eccentricity and on the total mass of the system. It does not depend on additional parameters that define the dynamics of the orbit at early inspiral, i.e., the value of the anomaly (left panel of Figure 2), and depends only weakly on the mass ratio of the binary. The right panel of Figure 2 shows that the $\mathcal{FF}$ tends to be larger for systems with larger total mass, since differences in GW phase evolution between eccentric and circularized waveforms have less cycles to accumulate. In contrast, systems with low total mass live longer and the discrepancies in the waveforms accumulate over time, leading to larger phase discrepancies, and hence lower $\mathcal{FF}$s. Based on these results, we conclude that using a template bank of circularized waveforms, and the placement bank algorithm described in Section II, is be sufficient to recover signals with $M_{\rm Total} \sim 2.4M_\odot\,(6M_\odot)$ emitted by BNS systems whose eccentricity at a fiducial GW frequency of 14Hz is $e \lesssim 0.02\,(0.05)$ with $\mathcal{FF} \gtrsim 0.95$.

We compare our findings with those reported in [21]. Figures 3-5 in [21] show the overlap between eccentric waveforms generated with the eccentric time-domain x-model and two different circular PN approximants, namely, time-domain TaylorT4 2PN and the frequency domain TaylorF2 at 3.5PN. The differences in the overlaps between the eccentric model and the circular PN approximants are caused by the different PN orders of the models used to generate the waveforms. Furthermore, as discussed in [37], PN waveforms that are perturbatively equivalent can have very different phase evolutions and asymptotic structure. These features are clearly exhibited in Figures 3-5 in [21]. Nonetheless, as pointed out in [21], maximization over the template bank gives rise to a significant gain in overlap. Having said that, we can compare our results with those computed in the context of initial LIGO in Figure 6 of [21]. Therein, it was found that initial LIGO may be able to efficiently detect BNS systems with $M_{\rm Total} \sim 2.4M_\odot\,(6M_\odot)$ with residual eccentricity $e \lesssim 0.05\,(0.08)$ at 40Hz. The authors in [21] also probed the effect of eccentricity in the context of advanced LIGO assuming the PSD given in Eq. (29) of [21] and using a low-frequency cut off of 10Hz. Given the demanding computational cost of such an analysis, they injected a few hundred signals to obtain a general picture of the efficacy with which Advanced LIGO could recover eccentric BNS signals. We have improved that initial analysis, and have carried out a large scale analysis assuming the expected sensitivity of advanced LIGO, a filter frequency of 15Hz, and an injection frequency of 14Hz. Taking into account the differences in low-frequency cut offs and detector noise spectrum, we notice a very good agreement between our results in the right-hand panel of Figure 2 and those reported in Figure 8 of [21]. We should point out, however, that we obtain more accurate results for low-mass BNSs because we have injected the eccentric signals at 14Hz.

IV. CONCLUSIONS

Compact binaries formed in core-collapsed globular clusters [13] or near supermassive black holes in galactic nuclei [19] may have high residual eccentricities when emitting gravitational radiation in LIGO band, however there is significant uncertainty on the rate of these events. An interesting prospect for advanced detectors is their ability to detect the GW radiation emitted by BNS systems in wide pre-supernovae orbits that are driven to coalescence in very short times when receiving appropriately oriented retrograde kicks. These type of events would be observed as a supernova followed by a extremely fast coalescence that takes the form of



FIG. 2. The left panel shows the Fitting Factor ($\mathcal{FF}$) as a function of the initial value of the eccentricity and the anomaly of the trajectory of the binary system. The right panel shows the correlation of the $\mathcal{FF}$ with the initial orbital eccentricity and the total mass of the binary system.

a second consecutive catastrophic event. Since both events may take place in the same location within months or years apart, these events may be effectively detected as double supernovae. Even though highly eccentric BNSs may not contribute significantly to the total number of BNSs that may be detected by advanced ground-based GW antennas, this study is timely and relevant to shed light on the selection bias that advanced searches of BNSs may incur by neglecting eccentricity.

We carried out a large scale Monte Carlo analysis to show that assuming that the expected noise curve of advanced LIGO is given by the Zero Detuned High Power PSD, BNS systems with total mass $\sim 2.4M_\odot\,(6M_\odot)$ and initial eccentricity $e \lesssim 0.02\,(0.05)$ at a fiducial GW frequency of 14Hz could be recovered with $\mathcal{FF} \gtrsim 0.95$ using a template bank of circularized waveforms. In order to detect and study the rate of eccentric BNS in aLIGO, a search specifically targeting these systems will need to be constructed. The construction of such a template matched filter search, as well as the extension of this study to higher mass systems [38, 39], is the subject of future work.

ACKNOWLEDGMENTS

This work was supported by NSF grants PHY-0847611 and PHY-0854812. DB and EH would also like to thank the Research Corporation for Science Advancement and the Cottrell Scholars program for support. The Monte Carlo simulations described in this paper were performed using the Syracuse University Gravitation and Relativity (SUGAR) cluster, which is supported by NSF grants PHY-1040231, PHY-0600953 and PHY-1104371. EH is pleased to thank Stefan Ballmer, Alessandra Buonanno, Peter Couvares, Jolien Creighton, Ryan Fisher, Chris Fryer, Ian Harry, Prayush Kumar, Alex Nitz, Bangalore Sathyaprakash, Peter Saulson and Matt West for useful discussions. We thank Ray Frey for carefully reading this manuscript and providing useful suggestions and criticisms. Part of this work was carried out at the Kavli Institute for Theoretical Physics at Santa Barbara University, which is supported in part by the National Science Foundation under Grant No. NSF PHY05-51164

---

[1] K. S. Thorne, in *Three hundred years of gravitation*, edited by S. Hawking and W. Israel (Cambridge University Press, 1987) pp. 330–458.
[2] C. S. Kochanek and T. Piran, Astrophys.J. **417**, L17 (1993), arXiv:astro-ph/9305015 [astro-ph].
[3] B. D. Metzger and E. Berger, Astrophys. J. **746**, 48 (2012).
[4] G. M. Harry and LIGO Scientific Collaboration, Classical and Quantum Gravity **27**, 084006 (2010).
[5] F. Acernese *et al.*, Classical and Quantum Gravity **25**, 184001 (2008).
[6] K. Somiya and the KAGRA Collaboration, Classical and Quantum Gravity **29**, 124007 (2012).
[7] J. Abadie *et al.*, Classical and Quantum Gravity **27**, 173001 (2010), arXiv:1003.2480 [astro-ph.HE].
[8] S. Osłowski, T. Bulik, D. Gondek-Rosińska, and K. Belczyński, *MNRAS* **413**, 461 (2011).
[9] I. Mandel and R. O'Shaughnessy, Classical and Quantum Gravity **27**, 114007 (2010).
[10] P. C. Peters and J. Mathews, *Phys. Rev.* D **131**, 435 (1963).
[11] V. Kalogera, R. Narayan, D. N. Spergel, and J. H. Taylor, Astrophys. J. **556**, 340 (2001).
[12] S. Barthelmy *et al.*, *Nature* **438**, 994 (2005).
[13] J. Grindlay, S. Portegies Zwart, and S. McMillan, Nature Physics **2**, 116 (2006), arXiv:astro-ph/0512654.
[14] E. Berger *et al.*, *Nature* **438**, 988 (2005).

[15] W. E. East and F. Pretorius, ArXiv e-prints (2012).

[16] B. A. Jacoby *et al.*, *Astrophys. J.* Lett **644**, L113 (2006).

[17] D. Guetta and T. Piran, *A&A* **453**, 823 (2006).

[18] R. M. O'Leary, B. Kocsis, and A. Loeb, *MNRAS* **395**, 2127 (2009).

[19] F. Antonini and H. B. Perets, Astrophys. J. **757**, 27 (2012).

[20] H. K. Chaurasia and M. Bailes, Astrophys. J. **632**, 1054 (2005).

[21] D. A. Brown and P. J. Zimmerman, Phys. Rev. D **81**, 024007 (2010).

[22] S. E. Thorsett and D. Chakrabarty, Astrophys. J. **512**, 288 (1999).

[23] D. Shoemaker, "Advanced LIGO anticipated sensitivity curves," (2010), `https://dcc.ligo.org/cgi-bin/DocDB/ShowDocument?docid=2974`.

[24] I. Hinder, F. Herrmann, P. Laguna, and D. Shoemaker, Phys. Rev. D **82**, 024033 (2010).

[25] B. Sathyaprakash and S. Dhurandhar, Phys.Rev. **D44**, 3819 (1991).

[26] R. Balasubramanian, B. Sathyaprakash, and S. Dhurandhar, Phys.Rev. **D53**, 3033 (1996).

[27] L. A. Wainstein and V. D. Zubakov, *Extraction of Signals from Noise* (Prentice-Hall, Englewood Cliffs, 1962).

[28] B. Allen, W. G. Anderson, P. R. Brady, D. A. Brown, and J. D. Creighton, Phys.Rev. **D85**, 122006 (2012).

[29] T. Apostolatos, Phys.Rev. **D52**, 605 (1995).

[30] B. J. Owen and B. S. Sathyaprakash, Phys. Rev. D **60**, 022002 (1999).

[31] S. Babak, R. Balasubramanian, D. Churches, T. Cokelaer, and B. Sathyaprakash, Class.Quant.Grav. **23**, 5477 (2006).

[32] A. Buonanno, Y.-b. Chen, and M. Vallisneri, Phys.Rev. **D67**, 104025 (2003).

[33] M. Boyle, D. A. Brown, L. E. Kidder, A. H. Mroue, H. P. Pfeiffer, *et al.*, Phys.Rev. **D76**, 124038 (2007).

[34] L. Blanchet, T. Damour, B. R. Iyer, C. M. Will, and A. G. Wiseman, Phys. Rev. Lett. **74**, 3515 (1995).

[35] L. Blanchet, T. Damour, G. Esposito-Farese, and B. R. Iyer, Phys. Rev. Lett. **93**, 091101 (2004).

[36] L. Blanchet, G. Faye, B. R. Iyer, and B. Joguet, Phys. Rev. **D71**, 129902 (2005).

[37] A. Buonanno, B. R. Iyer, E. Ochsner, Y. Pan, and B. S. Sathyaprakash, Phys. Rev. D **80**, 084043 (2009).

[38] E. A. Huerta and J. R. Gair, *Phys. Rev.* D **83**, 044020 (2011), arXiv:1009.1985 [gr-qc].

[39] E. A. Huerta and J. R. Gair, *Phys. Rev.* D **83**, 044021 (2011), arXiv:1011.0421 [gr-qc].